

Magnetic and Critical Properties of Alternating Spin Chain with $S = 1/2, 1$ in Magnetic Fields

Mitsuru Fujii¹, Satoshi Fujimoto² and Norio Kawakami^{1,3}

¹ *Department of Applied Physics, Osaka University, Suita, Osaka 565, Japan*

² *Department of Physics, Kyoto University, Kyoto 606, Japan*

³ *Department of Material and Life Science, Osaka University, Suita, Osaka 565, Japan*

(Received February 1, 2008)

We study an integrable spin chain with an alternating array of spins $S = 1/2, 1$ in external magnetic fields using the Bethe ansatz exact solution. The calculated magnetization possesses a cusp structure at a critical magnetic field $H = H_C$, at which the specific heat shows a divergence property. We also calculate finite-size corrections to the energy spectrum, and obtain the critical exponents of correlation functions with the use of conformal field theory (CFT). Low-energy properties of the model are described by two $c = 1$ $U(1)$ CFTs in $H < H_C$ and one $c = 1$ $U(1)$ CFT in $H > H_C$.

KEYWORDS: alternating spin chain, exact solution, conformal field theory

It is now well established that the antiferromagnetic quantum spin chains with half-integer spins have massless excitations, whereas those with integer spins have massive ones (Haldane gap), although the ground state of both cases are known to be singlet. Experimentally, another interesting class of quantum spin systems have been found, i.e. the spin systems with an alternating array of integer and half-integer spins both for ferromagnetic and antiferromagnetic cases¹. The systems found so far have the magnetically ordered ground state even for the antiferromagnetic case, for which quantum fluctuations may not play a crucial role. It is interesting to explore what happens if the magnetic order disappears, for example, in the presence of certain interactions beyond nearest neighbor ones. If such a system is realized, quantum fluctuations are expected to become important. This problem may thus provide us with a new interesting paradigm of quantum spin chains.

As a first step to address this problem of alternating spin chains with a singlet ground state, it may be interesting to study an integrable spin model^{2,3,4,5} whose ground state is singlet. In this paper, we study the integrable alternating spin model of $S = 1/2, 1$, with a special attention to its critical properties in magnetic fields. We systematically evaluate the magnetization, the specific heat and the critical exponents of correlation functions, and discuss their characteristic properties.

We consider a system of N spins $\frac{1}{2}\sigma_2, \frac{1}{2}\sigma_4, \dots, \frac{1}{2}\sigma_{2N}$ of spin- $\frac{1}{2}$ and N spins $S_1, S_3, \dots, S_{2N-1}$ of spin-1 in an external magnetic field H , the Hamiltonian for which is given by²

$$\mathcal{H} = \frac{1}{9} \sum_{n=1}^N \{ (2\sigma_{2n} \cdot S_{2n+1} + 1) (2\sigma_{2n+2} \cdot S_{2n+1} + 3) \\ + (2\sigma_{2n} \cdot S_{2n-1} + 1) [(\sigma_{2n-1} \cdot \sigma_{2n+1} + 1) (2\sigma_{2n} \cdot S_{2n+1} + 3) + (\sigma_{2n-1} \cdot \sigma_{2n+1} + 1) (2\sigma_{2n} \cdot S_{2n-1} + 1)] \}$$

where periodic boundary conditions are imposed: $\sigma_{2n} = \sigma_{2n+2N}$ and $S_{2n-1} = S_{2n-1+2N}$. Note that the Hamiltonian contains rather complicated next-nearest-neighbor interactions in addition to nearest-neighbor Heisenberg-type interactions, in order for the model to be integrable. The ground state of the model is known to be singlet^{2,3,4,5}, for which, as will be shown below, the antiferromagnetic correlation is dominant. In spite of its ugly appearance (1), we wish to note that this system may provide a starting model to discuss fundamental properties of alternating spin chains with quantum disordered (singlet) ground states. In the end of the paper, we shall give some discussions about what would be expected beyond this special model.

Let us first summarize the basic equations necessary for the following analysis. The Bethe equation obtained via the quantum inverse scattering method for (1) is

$$\left(\frac{\lambda_j - \frac{i}{2}}{\lambda_j + \frac{i}{2}} \right)^{\frac{L}{2}} \left(\frac{\lambda_j - i}{\lambda_j + i} \right)^{\frac{L}{2}} = - \prod_{l=1}^M \frac{\lambda_j - \lambda_l - i}{\lambda_j - \lambda_l + i}, \quad (2)$$

where $L = 2N$ is assumed to be a multiple of 4. Here we have taken the fully polarized state as a reference state, and denotes the number of “particles” by M , which are created by the spin-lowering operator. So, the z -component of the total spin is to be $S^z = 3N/2 - M$. One can see two kinds of phase factors reflecting the mixture of spin 1 and $1/2$ in the left-hand side of (2). The total energy is

$$E = \frac{7}{6}N - \sum_{j=1}^M \left(\frac{1}{\lambda_j^2 + \frac{1}{4}} + \frac{2}{\lambda_j^2 + 1} \right) - HS^z \quad (3)$$

and the total momentum is given by $P = \text{Im} \sum_{j=1}^M \log[(\lambda_j - \frac{i}{2})(\lambda_j - i)]$. So far, thermodynamic properties of the model have been investigated, revealing some properties distinct from those of the ordinary Heisenberg spin chain^{2,3,4,5}. Also, conformal properties have been discussed to some extent: the system at zero field is classified as

CFT of central charge $c = 2^{3,4)}$, and is conjectured to be described by level-1 $SU(2)$ Wess-Zumino-Witten models⁴⁾. In what follows, we systematically study the model in the presence of magnetic fields, and show that below a certain critical field H_C , low-energy properties of the model are described by two $c = 1$ $U(1)$ CFTs, while for $H > H_C$ by one $c = 1$ $U(1)$ CFT. Accordingly, the system shows quite different critical behaviors for low fields $H < H_C$ and high fields $H > H_C$.

We start with the calculation of the magnetization curve. To this end we need the Bethe equations in the thermodynamic limit. For large enough L , the ground state properties can be described by M_R real rapidities λ_j^R and 2-string complex rapidities λ_j^C ,

$$\lambda_j^\alpha = \lambda_j^C + \frac{1}{2}i(3 - 2\alpha) \quad (\alpha = 1, 2), \quad (4)$$

where λ_j^C ($j = 1, 2, \dots, M_C$) is a real number which labels the center of the 2-string^{2,3,4,5)}. The Bethe equation (2) is thus converted into integral equations for the density functions $\rho^R(\lambda^R)$ and $\rho^C(\lambda^C)$ for these rapidities

$$\begin{pmatrix} \rho^R \\ \rho^C \end{pmatrix} = \begin{pmatrix} \rho_0^R \\ \rho_0^C \end{pmatrix} - \begin{pmatrix} K_{RR} & K_{CR} \\ K_{RC} & K_{CC} \end{pmatrix} \otimes \begin{pmatrix} \rho^R \\ \rho^C \end{pmatrix}, \quad (5)$$

where the integral kernels are given by $K_{RR}(\lambda) = K_1(\lambda)$, $K_{CC}(\lambda) = \{2K_1(\lambda) + K_2(\lambda)\}$, $K_{CR}(\lambda) = K_{RC}(\lambda) = \{K_{1/2}(\lambda) + K_{3/2}(\lambda)\}$, respectively, in terms of the function $K_n(x) = (n/\pi)[n^2 + x^2]^{-1}$. The driving terms read

$$\begin{pmatrix} \rho_0^R \\ \rho_0^C \end{pmatrix} = \frac{1}{2} \begin{pmatrix} K_1 + K_{1/2} \\ K_{1/2} + K_1 + K_{3/2} \end{pmatrix}, \quad (6)$$

and the symbol $\otimes$ denotes the convolution integral in the finite range of $[\Lambda_\alpha^-, \Lambda_\alpha^+]$ respectively for the sectors of $\alpha = R, C$.

By calculating the above integral equations (5) numerically⁶⁾, we have computed the magnetization $\mathcal{M}(\equiv S^z)$ as a function of the magnetic field. The results are shown in Fig.1. We can see that the magnetization begins linearly in magnetic fields (finite spin susceptibility) since the ground state is singlet, and increases smoothly up to the critical field H_C , where we encounter a characteristic cusp structure⁷⁾. The existence of the cusp at $H = H_C$ indicates that the magnetization for $H < H_C$ consists of two contributions, i.e. $\mathcal{M} = \mathcal{M}_R + \mathcal{M}_C$, and one of them, $\mathcal{M}_C$, saturates at $H = H_C$. For $H > H_C$, the remaining massless spin excitations still contribute to the magnetization, and then at $H = 6$ the system is completely polarized. These characteristic properties directly reflect that the system consists of two kinds of different spins. The anomaly at $H = H_C$ also appears in the specific heat. We show the computed results for the coefficient γ of the T -linear specific heat in Fig.2. The divergence property for $H \rightarrow H_C$ implies that the velocity for one of spin excitations tends to zero when the system approaches the critical field H_C , since the specific heat coefficient γ is inversely proportional to the velocity. To see this point clearly, we have plotted two velocities

v_R and v_C in Fig.3. It is seen that v_C indeed vanishes at $H = H_C$. Therefore, we can say that there exist two species of spin excitations which behave differently in magnetic fields. They have the distinct pseudo-Fermi surfaces p_F^R and p_F^C , both of them decrease with the increase of magnetic fields. In particular, at $H = H_C$, p_F^C vanishes and the corresponding spin excitation becomes massive. In what follows we refer to these spin excitations as spinons in the R- and C-sectors respectively.

In order to further study critical properties of the model in magnetic fields, we now calculate the finite-size spectrum. Following a standard technique⁸⁾ we obtain the corrections to the energy

$$E = L\varepsilon_\infty(\pm\lambda_R^0, \pm\lambda_C^0) - \frac{\pi v_R}{6L} - \frac{\pi v_C}{6L} + \frac{2\pi v_R}{L}(\Delta_R^+ + \Delta_R^-) + \frac{2\pi v_C}{L}(\Delta_C^+ + \Delta_C^-). \quad (7)$$

where ε_∞ is the bulk energy density in the thermodynamic limit. Also, the finite-size corrections to the momentum are

$$P = \sum_{\alpha=R,C} [-2p_F^\alpha \Delta D_\alpha - \frac{2\pi}{L}(\Delta_\alpha^+ - \Delta_\alpha^-)] \quad (8)$$

where “the Fermi momenta” for two spinons are given by $p_F^R = \pi/4 - \pi\mathcal{M}_R/L$ and $p_F^C = \pi/2 - \pi\mathcal{M}_C/L$ with $\mathcal{M}_R = L/4 - M_R$ and $\mathcal{M}_C = L/2 - 2M_C$. Here ΔD_α labels an excitation across the two Fermi points, which carries the large momentum transfer $2p_F^\alpha \Delta D_\alpha$.

We note that both of the above expressions (7) and (8) completely fit in with the finite-size scaling formulae in CFT⁹⁾. Namely, the corrections $-\pi v_\alpha/(6L)$ to the ground state energy in (7) indicate that two kinds of spinons are described by two $c = 1$ CFTs. Also, the conformal dimensions $\Delta_\alpha^\pm$ which enter the excitation energy as well as the momentum are of desirable forms predicted by CFT,

$$\Delta_\alpha^\pm = \frac{1}{2} \left\{ \left(\frac{\xi_{\beta\beta} \Delta M_\alpha - \xi_{\alpha\beta} \Delta M_\beta}{2\det[\xi]} \right) \pm (\xi_{\alpha\alpha} \Delta D_\alpha + \xi_{\beta\alpha} \Delta D_\beta) \right\}^2 + n_\alpha^2 \quad (9)$$

for $\alpha = R, C$. Here the so-called dressed charge matrix $\xi_{\alpha\beta}$ ¹⁰⁾ is determined by the following integral equations,

$$\xi = I - K \otimes \xi, \quad (10)$$

where K is the 2×2 matrix of integral kernels which has appeared in eq.(5), and I is the identity matrix. The above dressed charge matrix determines the critical exponents of the present model in magnetic fields. Note that the conformal dimensions (9) both for R- and C-sectors have a typical form inherent in $c = 1$ Gaussian CFT with $U(1)$ symmetry, which has previously appeared in various contexts, for instance, in the analysis of interacting electron systems¹¹⁾. Recall here that for $H > H_C$, spinons of the C-sector become massive with vanishing v_C . Therefore we can see that low-energy properties of the present model in magnetic fields are described by two $c = 1$ CFTs for $0 < H < H_C$ and one $c = 1$ CFT for $H \geq H_C$.

Having specified the critical behavior in magnetic fields by CFT, we now read critical exponents of correlation functions from the conformal dimensions (9) by

suitably choosing the quantum numbers. For this purpose, we should note that the quantum numbers in (9) are subject to the selection rules

$$\Delta D_R = \frac{\Delta M_R}{2}, \quad \Delta D_C = \frac{\Delta M_C}{2} \pmod{1}, \quad (11)$$

which directly reflect that spinons can be regarded as interacting bosons.

For $H < H_C$ both of two kinds of spinons control the long-distance behavior of correlation functions, since v_R and v_C have finite values. Thus choosing the quantum numbers as

$$(\Delta M_R, \Delta M_C, \Delta D_R, \Delta D_C) = \begin{cases} (0, 0, 1, 0) \\ (0, 0, 0, 1) \end{cases}, \quad (12)$$

we find the spin-spin correlation function to have the asymptotic form,

$$\chi_z(x) \sim \frac{a_0}{x^2} + \frac{a_1 \cos(2p_F^R x)}{x^{\alpha_s^R}} + \frac{a_2 \cos(2p_F^C x)}{x^{\alpha_s^C}} \quad (13)$$

Note that the spin-spin correlation function has two oscillating terms, reflecting the presence of two-types of spinons. By inserting (12) into (9), and using the formula $\alpha = 2 \sum_{p=\pm} [\Delta_C^p + \Delta_R^p]$, we obtain critical exponents for the oscillating parts,

$$\alpha_s^R = 2(\xi_{RR}^2 + \xi_{RC}^2), \quad \alpha_s^C = 2(\xi_{CR}^2 + \xi_{CC}^2). \quad (14)$$

The numerical results for α_s^R and α_s^C are plotted as a function of the magnetic field in Fig.4. In weak magnetic fields, α_s^C represents how dominant the antiferromagnetic fluctuations are, because $p_F^C \sim \pi/2$. Since the value of α_s^C is close to unity in weak fields, we can see that the antiferromagnetic correlation is indeed dominant there, but is gradually suppressed with the increase of the magnetic field. For $H \geq H_C$, spinons of the C-sector become massive, and remaining massless spinons in the R-sector solely determine the critical behavior of the system. Accordingly, the third term of the right-hand side in eq.(13) vanishes. Also, the critical exponent α_s^R exhibits a discontinuity at $H = H_C$. This decrease in the critical exponent α_s^R may imply that the corresponding spin correlation may be enhanced to some extent in magnetic fields. Similar phenomena have been indeed observed in a different context where spin correlations are enhanced by decreasing the multiplicity of bands⁽¹²⁾. As seen from Fig.4, however, the corresponding exponent is larger than unity, so this fluctuation may not become dominant. We note here that a logarithmic correction should enter the correlation functions at $H = 0$ since the symmetry of the system is enhanced from $U(1)$ to $SU(2)$ at $H = 0$, although it does not appear in the present case with finite magnetic fields.

Here, following the arguments of Sachdev⁽¹³⁾, we would like to briefly discuss the temperature dependence of nuclear magnetic relaxation rates which may be an experimental probe to study low-energy properties of the system. Since the low-temperature behavior of the model is mainly determined by spinons mentioned above, we can obtain the leading temperature dependence of the nuclear spin-lattice relaxation rates $1/T_1$ and the nuclear

spin-spin relaxation rates $1/T_{2G}$ from eq.(13)⁽¹³⁾,

$$\frac{1}{T_1} \sim \frac{C_1}{T^{1-\alpha_s^R}} + \frac{C_2}{T^{1-\alpha_s^C}} + C_3 T, \quad (15)$$

$$\left(\frac{1}{T_{2G}} \right)^2 \sim \frac{D_1}{T^{3-2\alpha_s^R}} + \frac{D_2}{T^{3-2\alpha_s^C}}. \quad (16)$$

Combining these expressions with the numerical results for the critical exponents shown in Fig.4, we see that $1/T_1$ behaves like $\sim 1/T^{1-\alpha_s^C}$ at low temperatures for $H < H_C$. However for $H \geq H_C$ this term vanishes and the low temperature behavior of $1/T_1$ is determined by the term $\sim 1/T^{1-\alpha_s^R}$. The temperature dependence of $1/T_{2G}$ for $H < H_C$ is also governed by spinons in the C-sector and given by $1/T^{3/2-\alpha_s^C}$. For $H \geq H_C$ $1/T_{2G}$ behaves like $\sim 1/T^{3/2-\alpha_s^R}$.

We have been concerned with the rather special model with alternating spins in this paper. Before concluding the paper, we wish to briefly mention what would be expected if the system were away from the solvable point with coupling constants being changed. Unfortunately there have not been systematic attempts to this problem so far, so that we summarize here two possible disordered ground states predicted by the present analysis. (i) A possibility is that one of the $c = 1$ fields becomes massive in the presence of some relevant perturbations. In this case, we can see from the present results that low-field properties are classified by $c = 1$ CFT, and then the magnetization begins linearly. What is characteristic for this case is that a cusp structure in Fig.1 may not be observed. (ii) Another interesting possibility may be that both of two spinons become massive, and hence the system shows properties like a Haldane-gap system, for which there is no magnetization in low fields. These states, as well as the present massless state, should compete with the magnetically ordered state. In order to fully understand these problems, an extensive study with the aid of bosonization methods and numerical methods would be quite helpful. This issue is now under consideration.

In summary we have investigated magnetic and critical properties of an integrable alternating spin chain in external magnetic fields. The critical properties in magnetic fields are classified in terms of CFT. Although the present analysis is based on a rather special model, it would provide a plausible guideline to systematically explore characteristic properties of this new class of quantum spin chains.

-
- [1] G. T. Yee, J. M. Manriquez, D. A. Dixon, R. S. McLean, D. M. Groski, R. B. Flippen, K. S. Narayan, A. J. Epstein, and J. S. Miller: *Adv. Mater.* **3** (1991) 309; *Inorg. Chem.* **22** (1983) 2324; *ibid.* **26** (1987) 138.
 - [2] H. J. de Vega and F. Woyrnarovich: *J. Phys.* **A** **25** (1992) 4499.
 - [3] H. J. de Vega, L. Mezincescu, and R. I. Nepomechie: *Phys. Rev.* **B** **49** (1994) 13223; *Int. J. Mod. Phys.* **B** **8** (1994) 3473.
 - [4] S. R. Adams and M. J. Martins: *J. Phys.* **A** **26** (1993) L529.
 - [5] M. J. Martins: *J. Phys.* **A** **26** (1993) 7301.
 - [6] The cut-off parameters $\Lambda_\alpha^\pm$ are determined so as to minimize the free energy.
 - [7] We note that a similar cusp structure has been found for the

related spin model in which different spins are randomly distributed. See P. Schlottmann, Phys. Rev. B **49** (1994) 9202.

- [8] H. J. de Vega and F. Woynarovich: Nucl. Phys. **B 251** (1985) 439 (1985); F. Woynarovich, J. Phys. **A 22** (1989) 4243.
- [9] J.L. Cardy: Nucl. Phys. B **270**[FS16] (1986) 186; H. W. Bröte, J. L. Cardy and M. P. Nightingale: Phys. Rev. Lett. **56** (1986) 742; I. Affleck, Phys. Rev. Lett. **56** (1986) 746.
- [10] A.G. Izergin, V.E. Korepin and N.Y. Reshetikhin: J. Phys. **A22** (1989) 2615.
- [11] H.Frahm and V.E. Korepin: Phys. Rev. B. **42**, (1990) 10553; N.Kawakami and S.K. Yang: Phys. Rev. Lett. **65** (1990) 2309.
- [12] T. Itakura and N. Kawakami: J. Phys. Soc. Jpn. **64** (1995) 2321.
- [13] S. Sachdev: Phys. Rev. **B 50** (1994) 13006.

Fig. 1. Magnetization as a function of the magnetic field.

Fig. 2. Specific-heat coefficient as a function of the magnetic field.

Fig. 3. Velocities of spin excitations as a function of the magnetic field.

Fig. 4. Critical exponents α_s^R and α_s^C for oscillating sectors in the spin correlator as a function of the magnetic field.